

A New Method To Classify Flares Of UV Ceti Type Stars: Differences Between Slow And Fast Flares

H. A. Dal and S. Evren

Department Of Astronomy and Space Sciences, University of Ege,
Bornova, 35100 İzmir, Turkey

`ali.dal@ege.edu.tr`

Received _____; accepted _____

Not to appear in Nonlearned J., 45.

ABSTRACT

In this study, a new method is presented to classify flares derived from the photoelectric photometry of UV Ceti type stars. Using *Independent Samples t-Test*, the method is based on statistical analysis. The data used in the analyses were obtained from four flare stars observed between the years 2004 and 2007. Total number of flares obtained in the observations of AD Leo, EV Lac, EQ Peg and V1054 Oph is 321 in the standard Johnson U band. As a result, flare can be separated into two types as *slow* and *fast* depending on the ratio of *flare decay time* to *flare rise time*. The ratio is below the value 3.5 for all slow flares, while it is above 3.5 for all fast flares. Also, according to the *Independent Samples t-Test*, there are about 157 seconds difference between equivalent durations of slow and fast flares. In addition, there are significant differences between amplitudes and rise times of slow and fast flares.

Subject headings: methods: data analysis — methods: statistical — stars: flare — stars: individual(AD Leo, EV Lac, EQ Peg, V1054 Oph)

1. Introduction

Flares and flare processes are very hard worked subtitles of astrophysics. Lots of studies on flares have been carried out since the first flare was detected on the Sun by R.C. Carrington and R. Hodgson in 1 September 1959. Flare processes have not been perfectly understood yet (Benz & Güdel 2010). However, the researches indicate that the incidence of red dwarfs in our galaxy is 65%. Seventy-five per cent of them show flare activity, these stars are known as UV Ceti type stars (Rodonó 1986). In this respect, it will be easier to understand the evolution of red dwarfs if the flare processes are well known. This is because flare activity dramatically affects the evolutions of the red dwarfs. In this respect, an attempt to classify flares by considering shapes of flare light variations observed in the UV Ceti type stars, flares have been tried to classify. It is believed that classification of flares makes the flares and flare processes more intelligible.

Flares of the UV Ceti type stars were first detected in 1939 (Van Maanen 1940). Discovering the flare stars with high flare frequency such as UV Cet, YZ CMi, EV Lac, AD Leo and EQ Peg, the detected flare numbers and the variety of flare light variations increased. The light variation of each flare is almost different from each other. In the first place, it is seen that there are lots of shapes for flare light variations (Moffett 1974; Gershberg 2005). On the other hand, when large numbers of flares are examined, it is seen that there are only two main shapes for flare light variations. One of them is called the fast flare. Fast flares have higher energy and the shapes of their light variations are similar to the shapes of solar hard X-ray flares. The second type flares are called slow flares. Unlike fast flares, slow flares exhibit lower energy. The rise times of slow flares are almost equal to their decay times.

The terms of fast and slow flares were used for the first time in 1960's in astrophysics. If the rise time of a flare is below 30 minutes, Haro & Parsamian (1969) called that

flare *a fast flare*. If its rise time is above 30 minutes, they called it *a slow flare*. Like Haro & Parsamian (1969), considering the shapes of flare light variations, Osawa et al. (1968) described two types of flares. However, Oskanian (1969) separated flares into four classes. Like Oskanian (1969), considering only light variation shapes of the flares, Moffett (1974) directly classified flares such as classical, complex, slow, and flare event. On the other hand, Kunkel had asserted another idea in his PhD thesis in 1967 (Gershberg 2005). According to Kunkel, the observed flare light variations must be some combinations of some slow and fast flares. According to this idea, there are two main flare types. The complex flares mentioned by Moffett (1974) are actually a combination of some fast and slow flares. And also, both slow flares and flare events mentioned by Moffett (1974) can be classified as the same type flares.

Gurzadyan (1988) described two flare types to model the flare light curves. Gurzadyan (1988) indicated that thermal processes are dominated in the processes of slow flares. And these flares are ninety-five per cent of all flares observed in UV Ceti type stars. The non-thermal processes are dominated in the processes of fast flares, which are all the other flares. According to Gurzadyan (1988), there is a large energy difference between these two types of flares.

In this study, we introduce a new statistical method for classifying flares. Using statistical *Independent Samples t-Test* (hereafter *t-Test*) analysis, this method is based on the distribution of flare equivalent durations versus flare rise times. Considering the studies in Osawa et al. (1968), Kunkels PhD thesis (Gershberg 2005) and Gurzadyan (1988), we assume that there are two flare types such as fast and slow flares types. We classify flares into two types and demonstrate the similarities and differences between these two types of flares. In respect of these analyses, we discuss the results obtained from analyses of 321 flares detected in U band observations of flare stars AD Leo, EV Lac, EQ Peg and V1054

Oph between 2004 and 2007. The programme stars were selected for this study due to their high flare frequencies (Moffett 1974). The flare data obtained in this study are useful for such analysis. This is because the data were obtained with systematical observations, using the same method.

The flare activity of AD Leo was discovered by Gordon & Kron (1949) for the first time. The star is a red dwarf and a member of The Castor Moving Group, whose age is about 200 million years (Montes et al. 2001). Crespo-Chac et al. (2006) found that the flare frequency of AD Leo was $0.71 h^{-1}$. The variation of the flare frequency was investigated by Ishida et al. (1991). They mentioned that there is no variation in the flare frequency of AD Leo. The other star in this study is EV Lac, which is one of the well known UV Ceti stars. According to the spatial velocities, EV Lac seems to be a member of 300 million years old Ursa Major Group (Montes et al. 2001). It has been known since 1950 that EV Lac shows flares (Lippincott 1952; Van de Kamp 1953). The largest observed flare amplitude is 6.4 mag in U band observations of EV Lac. Andrews (1982) detected 50 flares in U and B band observations of EV Lac. The author indicated that about 42 flares of 50 flares were found in some groups, which were occurred in every 5-6 days. The seasonal flare frequencies were computed from 1972 to 1981 and these frequencies were compared with the seasonal averages of B band magnitudes. According to this comparison, it was found that the activity cycle is about 5 years for EV Lac (Mavridis & Avgoloupis 1986). On the other hand, Ishida et al. (1991) indicated that there was no flare frequency variation from 1971 to 1988. In another study, Leto et al. (1997) showed that flare frequencies of EV Lac increased from 1968 to 1977. EQ Peg is another active flare star, whose flare activity was discovered by Roques (1954). EQ Peg is classified as a metal-rich star and it is a member of the young disk population in the galaxy (Veeder 1974; Fleming et al. 1995). EQ Peg is a visual binary (Wilson 1954). Both of the components are a flare star (Pettersen et al. 1983). Angular distance between components is given as a value between $3''.5$ and $5''.2$

(Haisch et al. 1987; Robrade et al. 2004). One of the components is 10.4 mag and the other is 12.6 mag in V band (Kukarin 1969). Observations show that flares of EQ Peg generally come from the fainter component (Fossi et al. 1995). Rodonó (1978) proved that 65% of the flares come from faint component and about 35% from the brighter component. The fourth star in this study is V1054 Oph, whose flare activity was discovered by Eggen (1965). V1054 Oph (= Wolf 630ABab, Gliese 644ABab) is a member of Wolf star group (Joy 1947; Joy & Abt 1974). Wolf 630ABab, Wolf 629AB (= Gliese 643AB) and VB8 (= Gliese 644C), are the members of the main triplet system, whose scheme is demonstrated in Fig.1 given in the paper of Pettersen et al. (1984). Wolf 630 and Wolf 629 are visual binary and they are separated $72''$ from each other. Wolf 630AB is a close visual binary in itself. Wolf 629AB is a spectroscopic binary. B component of Wolf 629AB seems to be a spectroscopic binary. VB8 is $220''$ far away from the other components. There is an angular distance about $0''.218$ between A and B components of Wolf 630 (Joy 1947; Joy & Abt 1974). The masses were derived for each components of Wolf 630ABab by Mazeh et al. (2001). The author showed that the masses are $0.41 M_{\odot}$ for Wolf 629A, $0.336 M_{\odot}$ for Wolf 630Ba and $0.304 M_{\odot}$ for Wolf 630Bb. In addition, Mazeh et al. (2001) demonstrated that the age of the system is about 5 Gyr.

2. Observations and Analyses

2.1. Observations

The observations were acquired with a High-Speed Three Channel Photometer attached to a 48 cm Cassegrain type telescope at Ege University Observatory. Using a tracking star set in a second channel of photometer, the observations were continued in standard Johnson U band with the exposure time between 2 and 10 seconds. The basic parameters of all program stars and their comparisons are given in Table 1. The parameters given in the

table are star name, magnitude in V band, B-V colour index, spectral type, distance (pc) and bolometric luminosity ($\text{Log}L_{bol}$, ergs s^{-1}). The magnitudes and colour indexes were obtained in this study. Considering B-V colour indexes, the spectral types were taken from Tokunaga (2000). The distances and bolometric luminosities were taken from Fossi et al. (1995) and Gershberg et al. (1999).

Although the program and comparison stars are so close on the sky, differential extinction corrections were applied. The extinction coefficients were obtained from the observations of the comparison stars on each night. Moreover, the comparison stars were observed with the standard stars in their vicinity and the reduced differential magnitudes, in the sense variable minus comparison, were transformed to the standard system using procedures outlined in Hardie (1962). The standard stars are listed in the catalogues of Landolt (1983, 1992). Heliocentric corrections were applied to the times of the observations. The standard deviation of observation points acquired in standard Johnson U band is about $0^m.015$ on each night. Observational reports of all program stars are given in the Table 2. It is seen that there is no variation of differential magnitudes in the sense comparison minus check stars.

Gershberg (1972) developed a method for calculating flare energies. Flare equivalent durations and energies were calculated with using Equation (1) and (2) of this method.

$$P = \int [(I_{flare} - I_0) / I_0] dt \quad (1)$$

In Equation (1), I_0 is the intensity of the star in quiescent level. I_{flare} is the intensity during flare.

$$E = P \times L \quad (2)$$

where E is the flare energy; P is the flare equivalent duration; L is the luminosity of the stars in quiescent level in the Johnson U band.

HJD of flare maximum times, flare rise and decay times, amplitudes of flares and flare equivalent durations were calculated for each flare. Brightness of a star without a flare was taken as a quiescent level of brightness of this star on each night. Considering this level, all flare parameters were calculated for each night. It was seen that some flares have a few peaks. In this case, flare maximum time and amplitudes were calculated from the first highest peak. Instead of flare energies, flare equivalent durations were used for all statistical analyses. This is because of the luminosity term in Equation (2). The luminosities of stars from different spectral types have great differences. Although the equivalent durations of two flares obtained from two stars in different spectral types are the same, calculated energies of these flares are different due to different luminosities of these spectral types. Therefore, we could not use these flare energies in the same analyses. On the other hand, flare equivalent duration depends just on flare power. Another reason of using equivalent duration is that the given distances of the same star in different studies are quite different. Therefore, the calculated luminosities become different because of these different distances.

All the calculated parameters of flares are given in Table 3. The given parameters in columns are star name, observation date, HJD of flare maximum, flare total time (s), decay time (s), equivalent duration (s), energy (ergs), flare amplitude (mag) and flare types. In the last column, it is given whether the flare was used in the analyses, or not.

When the observed flares are examined, it is seen that almost each flare has a distinctive light variation shape (Figures 1, 2, 3, and 4). In these figures, horizontal dashed lines

represent the level of quiescent brightness. The flare seen in Figure 1 is a fast flare. This flare type occurs frequently in UV Ceti type stars. On the other hand, the first flare seen in Figure 2 is a compact flare. This flare type among the others is the hardest flare type to classify. This flare type must be a combination of two flares. The observed flares, whose light variations are similar to the first flare in Figure 2, were not used in the analyses. The second flare seen in Figure 2 is a fast flare. The flare seen in Figures 3 is a powerful flare, but its light variation was not completed because we did not carry on observation until the flare completely decreased to quiescent phase due to the Sun rising. If the light variation of a flare was not completed like this one, the flare was not used in the analyses. The flare seen in Figure 4 seems to be very different from previous flares. Moffett (1974) called flares like this one as the flare events. They are called slow flares in some other studies. In this study, we called them as the slow flares.

2.2. Analyses

The impulsive phase of a flare is the time interval where sudden-high energy occurs. On the other hand, the mean phase is other part of the flare, where the energy is emitted to all space (Gurzadyan 1988; Benz & Güdel 2010). Moreover, rate of brightness increase for fast flares is clearly higher than that of slow flares (Gurzadyan 1988; Gershberg 2005). Moreover, Gurzadyan (1988) stated that, the energies of fast flares are always higher than slow flare energies.

According to this approach, we examined all flare data and we saw that the equivalent durations of some flares are different, while their rise times are the same. For example, the rise time of 22 flares is 15 second. The equivalent durations of eight flares among them are very high, while the equivalent durations of other 14 flares are dramatically low. It was seen in 30 different rise times. It means that there are least two flares in 30 different rise

times and their equivalent durations are different from each other. 140 flares were chosen in total. These 140 flares used in the analyses are specified in the last column of Table 3. Considering also their light variation shapes, we separated flares into two groups as flares with high energy and low energy. It was found that 61 flares have high energy, while 79 flares have low energy. Considering light variations, it was seen that 61 flares are fast flare, and the others are slow flare. For each one of 30 rise times, the averages of the equivalent durations were computed separately for the flares with high energy and low energy.

The most suitable statistical test for these data is the *t-Test* to determine the difference between the equivalent durations of two groups. This is because the *t-Test* examines whether there is any statistical difference between independent variable of two groups, or not (Wall & Jenkins 2003; Dawson & Trapp 2004). In this study, the flare rise times were taken as a dependent variable, while flare equivalent durations were taken as an independent variable. In the analyses, the SPSS V17.0 software was used (Green et al. 1999). The average of equivalent durations in the logarithmic scale for 79 slow flares was calculated and found to be 1.348 ± 0.092 . And it was found to be 2.255 ± 0.126 for 61 fast flares. This shows that there is a difference about 0.907 between average equivalent durations in the logarithmic scale. The *p-probability value* (hereafter *p-value*) was computed to test the results of the *t-Test* and it was found as *p-value* < 0.0001 . It means that the result is statistically acceptable. All the results of the analyses are given in Table 4.

In the next step, we compared their distributions, the distribution of the equivalent durations versus flare rise times. The best fits for distributions were searched. Using GraphPad Prism V5.02 software (Motulsky 2007), regression calculations showed that the best fits of distributions seen in Figure 5 were linear functions given by Equations (3) and (4).

$$\text{Log}(P_u) = 1.109 \times \text{Log}(T_r) - 0.581 \quad (3)$$

$$\text{Log}(P_u) = 1.227 \times \text{Log}(T_r) + 0.122 \quad (4)$$

It was tested whether these linear functions belong to two independent distributions, or not. In this point, the slopes of linear functions were principally examined. As it can be seen in Table 4, the slope of the linear function is 1.109 ± 0.127 for slow flares, while it is 1.227 ± 0.243 for fast flares. This shows that the increasing of equivalent durations versus flare rise times for both fast and slow flares are parallel. When the probability, *p-value*, was calculated to say whether it is statistically significant, it was found that *p-value* = 0.670. This value indicates that there is no significant difference between the slopes of fits.

Finally, the *y-intercept* values were calculated and compared for two linear fits. While this value is -0.581 for the slow flares, it is 0.122 for the fast flares in the logarithmic scale. It means that there is a difference about 0.703 between these values in the logarithmic scale. When the probability value was calculated for *y-intercept* values to say whether there is a statistically significant difference, it was found that *p-value* < 0.0001. This result indicates that the difference between two *y-intercept* values is clearly important.

Some other differences like ones demonstrated by *t-Test* are directly seen in the graphics. For example, the lengths of flare rise times for both type flares can be compared in Figure 6. While the lengths of rise times for slow flares can reach to 1400 seconds, they are not longer than 400 seconds for fast flares.

The comparison of another parameter is given in Figure 7. The flare amplitudes are seen in this figure. As it can be seen, while the amplitudes of fast flares can reach to 4.0 mag, the amplitudes of slow flares can exceed 1.0 mag.

61 fast and 79 slow flares were chosen among 321 flares observed in this study. The ratios of flare decay time to flare rise time were computed for both 61 fast and 79 slow flares. As a result, it is seen that the ratio is below the value of 3.5 for each one of 79 slow flares. On the other hand, the ratio is above the value of 3.5 for each fast flares. The value of 3.5 is considered as a limit for these two type flares. Considering the ratio of 3.5, other 181 flares of 321 flares were separated as slow and fast flares. When the results obtained from analyses of 140 flares were rechecked for 321 flares, it was seen that the results are the same with the previous ones.

3. Results and Discussion

We observed 321 flares in U band observations of AD Leo, EV Lac, EQ Peg and V1054 Oph. Examining 321 flares, 61 fast and 79 slow flares were identified for analyses. The *t-Test* was used as an analysis method. Flare rise times were accepted as dependent variables, while flare equivalent durations were taken as independent variables. The results obtained from the *t-Test* analyses of the data show that there are distinctive differences between two flare types. These differences are important properties because the models of white light flares observed in photoelectric photometry must support these properties to explain both flare types.

The distributions of the equivalent durations were represented by linear fits given by Equations (3) and (4) for these flare types. The slope of linear fit is 1.109 for slow flares, which are low energy flares. And, it is 1.227 for fast flares, which are high energy flares. The values are almost close to each other. It seems that the equivalent durations versus rise times increase in similar ways.

In the case of UV Ceti stars, when flare models are considered, it is seen that there are

two main energy sources for flares (Gurzadyan 1988; Benz & Güdel 2010). These depend on the thermal and non-thermal processes (Gurzadyan 1988). Flares with small amplitude are generally the flares with low energy. The thermal processes are commonly dominant for these type flares. On the other hand, the flares, which have sudden rapid increases, are more energetic events. Non-thermal processes are dominant for this type. And thus, there is an energy difference between these two types of flares (Gurzadyan 1988). When the averages of equivalent durations for two type flares were computed in logarithmic scale, it was found that the average of equivalent durations is 1.348 for slow flares and it is 2.255 for fast flares. The difference of 0.907 between these values in logarithmic scale is equal to 157.603 second difference between the equivalent durations. As it can be seen from Equation (2), this difference between average equivalent durations affects the energies in the same way. Therefore, there is 157.603 times difference between energies of these two type flare. This difference must be the difference mentioned by Gurzadyan (1988).

The slopes of linear fits are almost close. On the other hand, if the *y-intercept* values of the linear fits are compared, it is seen that there is 0.703 times difference in logarithmic scale, while there is 0.907 times difference between general averages. Considering also Figure 5, it is seen that equivalent durations of fast flares can increase more than slow flare equivalent durations towards the long rise times. Some other effects should be involved in the fast flare process for long rise times. These effects can make fast flares more powerful than they are.

When the lengths of rise times for both flare types are compared, it is seen that there is a difference between them. The lengths of rise times can reach to 1400 seconds for slow flares, but are not longer than 400 seconds for fast flares. In addition, when the flare amplitudes are examined for both type flares, an adverse difference is seen according to rise times. While the amplitudes of slow flares reach to 1.0 mag at most, the amplitudes of fast

flares can exceed 4.0 mag.

Finally, when the ratios of flare decay times to flare rise times are computed for two flares types, the ratios never exceed the value of 3.5 for all slow flares. On the other hand, the ratios are always above the value of 3.5 for fast flares. It means that if decay time of a flare is 3.5 times longer than its rise time at least, the flare is a fast flare. If not, the flare is a slow flare. Therefore, the type of an observed flare can be determined by considering this value of the ratio. In the studies like Osawa et al. (1968), Oskanian (1969), Haro & Parsamian (1969) and Moffett (1974), considering directly the shapes of flare light variations, the flares have been classified into two types as fast and slow flares. For instance, according to Haro & Parsamian (1969), if the rise time of a flare is above 30 minutes, the flare is slow flare. If not, it is a fast flare. However it is shown in this study that there are some fast flares, whose rise times are longer than the rise times of some slow flares. This is clearly seen from Table 3. This case indicates that a classification by considering only the rise time may not be right. Nevertheless, Moffett (1974) separated flares into more than two groups such as classic, complex, spike and flare events. On the other hand, according to our results of *t-Test* analyses, neither only one parameter nor the shape of the light variation was enough to classify a flare. The flare equivalent durations and also one more parameter should be taken into consideration in order to make such a classification.

The values 3.5, the ratio of flare decay times to flare rise times, can give an idea about the rate of energy emitting in a flare process. The rise times of flares are some limits for each type. Maximum flare rise time is about 400 seconds for fast flares, while it can reach the values over 1400 seconds for slow flares. However, the decay times can take any duration without any limited values. Consequently, the ratio of flare decay times to flare rise times depends on rise time more than decay times. In the case of rise time, the difference between

two type flares must be caused by whether the flare processes are thermal or non-thermal. We computed the duration as a rise time from the phase in which the brightness increases. Increasing of the brightness is caused by increasing the temperature of some region on the surface of the star. The flare rise time is an indicator of heating this region on the surface. Therefore, the ratio of flare decay times to flare rise times, so the values of 3.5, must be a critical value between thermal or non-thermal processes.

As it is seen from the models of Gurzadyan (1988), the differences between flare durations and flare amplitudes are seen between two flare types derived from observed flares in this study. The difference between amplitudes of slow and fast flares was given by Equation (22) in the paper of Gurzadyan (1988). In the case of flare amplitude, the result obtained in this study is in agreement with this equation.

Providing that the value 3.5 is a limit ratio for flare types, fast flare rate is 63% of all 321 flares observed in this study, while slow flare rate is 37%. It means that one of every three flares is a fast flare, while two of them are slow flares. This result diverges from what Gurzadyan (1988) stated. According to Gurzadyan (1988), slow flares with low energies and low amplitudes are 95% of all flares. The remainder are fast flares. When looking individually over each star, the rate of flare types is changing from star to star. Detected flare number of AD Leo is 110 as it can be seen from Table 2. Slow flare rate of AD Leo flares is 78%, while the rate of fast flares is 22%. Detected flare number is 98 in observation of EV Lac and 40 in observation of V1054 Oph. Slow flare rates of both stars are 75%, while fast flare rates are 25%. EQ Peg flare number is 78. Slow flare rate of them is 63%, and fast flare rate is 37%.

In this study, one of the remarkable points is the correlation coefficients of linear fits. As it is seen in Table 4, the correlation coefficient is 0.732 for linear fit of slow flare type and 0.476 for fast flares. Although the correlation coefficient of the linear fit to the

distribution of equivalent durations versus rise times is in an acceptable level for slow flares, it is relatively lower for fast flares. Regression calculations show that the best fits are linear for the distribution of equivalent durations versus rise times in logarithmic scales. The correlation coefficients of other fits are not higher than linear correlation coefficients. Especially, the correlation coefficient is lower for the fast flares due to the distribution of their equivalent durations. As it is seen from Figure 5, the equivalent durations of fast flares can take values in a wide range towards the longer rise times. This must be owing to the same reason of differences between *y-intercept* values and the mean averages of equivalent durations of two flare types. As it is discussed above, while the slopes of the fits are nearly close to each other, there is a considerable difference between *y-intercept* values and mean average of equivalent duration for two flare types. Consequently, all these deviations are seen in fast flares. The magnetic reconnection is dominant in this type of flares. A parameter in magnetic reconnection process causes some fast flares to be more powerful than the expected values. Eventually, some fast flares are more powerful than they are, while some of them are at expected energy levels. On the other hand, this parameter in magnetic reconnection process is not dominated in slow flare processes. And so, distribution of their equivalent durations is not scattered. This must be why the correlation coefficient of the fit is relatively higher for slow flares.

In this classification method, the complex flares are an exceptional case. These flares must be composed of some different flares. The complex flare should be separated into component flares before classification. If the fast and slow flares can be modelled, using these models, the complex flares can be decomposed into component flares.

In conclusion, some parameters can be computed from flares observed in photoelectric photometry. And, if the behaviours between these parameters can be analysed by suitable methods, the flare types can be determined. In this study, we analysed the distributions

of equivalent durations versus flare rise time by the statistical analysis method, *t-Test*. Finally, it is seen that using the ratios of flare decay times to flare rise times, flares can be classified. Thus, flares are classified into two types as fast and slow flares. It is seen that there are considerable differences between these two types of the flares. The differences and the similarities between flare types are important to understand the flare processes. This gives new ideas to model white light flares of UV Ceti stars.

Acknowledgments

The authors acknowledge generous allotments of observing time at Ege University Observatory. We wish to thank both Dr. Hayal Boyacıoğlu, who gave us important suggestions about statistical analyses, and Prof. Dr. M. Can Akan, who gave us valuable suggestions, which improved the language of the manuscript. We also thank the referee for useful comments which have contributed to improve the manuscript. We finally wish to thank the Ege University Research Found Council for supporting this work through grant Nr. 2005/FEN/051.

REFERENCES

- Andrews, A. D., 1982, IBVS, 2254, 1
- Benz, A. O. & Güdel, M., 2010, ARA&A, 48, (in press)
- Crespo-Chacón, I., Montes, D., García-Alvarez, D., Fernández-Figueroa, M. J., López-Santiago, J., Foing, B. H., 2006, A&A, 452, 987
- Dawson, B. & Trapp, R. G., 2004, "In Basic and Clinical Biostatistics", The McGraw-Hill Companies Inc. Press, USA, p.61, p.134, p.245
- Eggen, O. J., 1965, Obs, 85, 191
- Fleming, Th. A., Schmitt, J. H. M. M., Giampapa, M. S., 1995, ApJ, 450, 401
- Fossi, B. C. M., Landini, M., Fruscione, A., & Dupuis, J., 1995, ApJ, 449, 376
- Gershberg, R. E., 1972, Astrophys. Space Sci. 19, 75
- Gershberg, R. E., Katsova, M. M., Lovkaya, M. N., Terebizh, A. V., Shakhovskaya, N. I., 1999, A&AS, 139, 555
- Gershberg, R. E., 2005, "Solar-Type Activity in Main-Sequence Stars", Springer Berlin Heidelberg, New York, p.53, p.191, p.192, p.194, p.211, p.325, p.360
- Gordon, K. C., & Kron, G. E., 1949, PASP, 61,210
- Green, S. B., Salkind, N. J., Akey, T. M., 1999, "Using SPSS for Windows: Analyzing and Understanding Data", Upper Saddle River, N.J. ; London : Prentice Hall Press, P.50
- Gurzadyan, G. A., 1988, ApJ, 332, 183
- Haisch, B. M., Butler, C. J., Doyle, J. G., & Rodonó, M., 1987, A&A, 181, 96

- Hardie R.H., 1962, "in Astronomical Techniques", ed.W.A.Hiltner (Chicago: Univ. Chicago Press), 178
- Haro, G., & Parsamian, E., 1969, BOTT, 5, 45
- Ishida, K., Ichimura, K., Shimizu, Y., Mahasenaputra, 1991, Ap&SS, 182, 227
- Joy, A. H., 1947, ApJ, 105, 96
- Joy, A. H. & Abt, H. A., 1974, ApJS, 28, 1
- Kukarin, B. V., 1969, "in General Catalogue of Variable Stars", 3d ed., Moscow Sternberg Astronomical Institute
- Landolt, A. U., 1983, AJ, 88, 439
- Landolt, A. U., 1992, AJ, 104, 340
- Leto, G., Pagano, I., Buemi, C. S., Rodonó, M., 1997, A&A, 327, 1114
- Lippincott, S.L., 1952, ApJ, 115, 582
- Mavridis, L. N. & Avgoloupis, S., 1986, A&A, 154, 171
- Mazeh, T., Latham, D. W., Goldberg, E., Torres, G., Stefanik, R. P., Henry, T.J., Zucker, S., Gnat, O., Ofek, E. O., 2001, MNRAS, 325, 343
- Moffett, T. J., 1974, ApJS, 29, 1
- Montes, D., López-Santiago, J., Gálvez, M. C., Fernández-Figueroa, M. J., De Castro, E., Cornide, M., 2001, MNRAS, 328, 45
- Motulsky, H., 2007, "In GraphPad Prism 5: Statistics Guide", GraphPad Software Inc. Press, San Diego CA, p.94, p.133

- Osawa, K., Ichimura, K., Noguchi, T., & Watanabe, E., 1968, Tokyo Astron. Bull., No 180
- Oskanian, V. S., 1969, In: L. Detre (ed). "Non-Periodic Phenomena in Variable Stars",
Proc. AUI Coll. No 4. Academic Press, Budapest. p.131
- Pettersen, B. R., Kern, G. A. & Evans, D. S., 1983, A&A, 123, 184
- Pettersen, B. R., Coleman, L. A., Evans, D. S., 1984, ApJ, 282, 214
- Robrade, J., Ness, J. U., and Schmitt, J. H. M. M., 2004, A&A, 413, 317
- Rodonó, M., 1978, A&A, 66, 175
- Rodonó, M., 1986, NASSP, 492, 409
- Roques, P. E., 1954, PASP, 66, 256
- Tokunaga A. T., 2000, "Allen's Astrophysical Quantities", Fouth Edition, ed. A.N.Cox
(Springer), p.143, p.479
- Van de Kamp, P., 1953, PASP, 65, 73
- Van Maanen, A., 1940, ApJ, 91, 503
- Veeder, G. J., 1974, AJ, 79, 702V
- Wall, J. W. & Jenkins, C. R., 2003, "In Practical Statistics For Astronomers", Cambridge
University Press, p.79
- Wilson, R. H., Jr., 1954, AJ, 59, 132




Fig. 1.— A flare light curve sample for fast flares obtained from U band observation of V1054 Oph in 14 June 2004.




Fig. 2.— A flare light curve sample for fast flares obtained from U band observation of EV Lac in 10 August 2004.




Fig. 3.— A flare light curve sample for more powerful flares obtained from U band observation of AD Leo in 6 February 2007.




Fig. 4.— A flare light curve sample for slow flares obtained from U band observation of EQ Peg in 16 September 2004.




Fig. 5.— The distributions for the mean averages of the equivalent durations ($\text{Log}(P_u)$) versus flare rise times ($\text{Log}(T_r)$) in logarithmic scale. In the figure, open circles represent slow flares, while filled circles show the fast flares. And the lines represent fits given equations (3) and (4).




Fig. 6.— The distributions of the equivalent durations ($\text{Log}(P_u)$) in logarithmic scale versus flare rise times (T_r) for all 321 flares detected in observations of program stars. In the figure, open circles represent slow flares, while filled circles show the fast flares.




Fig. 7.— The distributions of flare amplitudes versus flare rise times ($\text{Log}(T_r)$) in logarithmic scale for all 321 flares detected in observations of program stars. In the figure, open circles represent slow flares, while filled circles show the fast flares.

Table 1: Basic parameters for the targets studied and their comparison (C1) and check (C2) stars.

Stars	V (mag)	B-V (mag)	Spectral Type	Distance (pc)	$LogL_{bol}$
AD Leo	9.388	1.498	M3	4.90	31.87
C1 = HD 89772	8.967	1.246	K6-K7	-	-
C2 = HD 89471	7.778	1.342	K8	-	-
EV Lac	10.313	1.554	M3	5.00	31.72
C1 = HD 215576	9.227	1.197	K6	-	-
C2 = HD 215488	10.037	0.881	K1-K2	-	-
EQ Peg	10.170	1.574	M3-M4	6.58	31.42
C1 = SAO 108666	9.598	0.745	G8	-	-
C2 = SAO 91312	9.050	1.040	K3-K4	-	-
V1054 Oph	8.996	1.552	M3	5.70	31.93
C1 = HD 152678	7.976	1.549	M3	-	-
C2 = SAO 141448	9.978	0.805	K0	-	-

Table 2: Observational reports of the each program star for each observing season.

Stars	Year	HJD (+2400000)	Filter	Observation Number	Observation Time (hour)	U Filter Flare Number
AD Leo	2005	53377 - 53514	U	12	36.35	39
	2006	53717 - 53831	U	15	37.48	54
	2007	54048 - 54248	U	8	20.80	17
EV Lac	2004	53197 - 53257	U	17	47.62	31
	2005	53554 - 53606	U	9	26.65	32
	2006	53940 - 53996	U	16	44.66	35
EQ Peg	2004	53236 - 53335	U	13	64.42	38
	2005	53621 - 53686	U	10	35.84	35
V1054 Oph	2004	53136 - 53202	U	19	42.64	14
	2005	53502 - 53564	U	10	33.13	26

Table 3. All the parameters were computed from observed flares. From the first column to the last, star name, the date of observation, HJD of flare maximum moment, flare total time (sec), decay time (sec), equivalent duration (sec), flare energy (erg), flare amplitude (mag) and flare type are given, respectively. And in the last column, it is specified whether the flare was used in the analyses, or not.

Stars	Observation Date	HJD For Maximum of Flare	Total Time (sec)	Decay Time (sec)	Equivalent Duration (sec)	Flare Energy (ergs)	Amplitude In U Band (mag)	Flare Type	Used In Analyses
AD Leo	06.01.2005	53377.50660	60	36	5.1690	6.8480E+30	0.348	Slow	Yes
AD Leo	06.01.2005	53377.59771	684	660	94.5297	1.2523E+32	0.723	Fast	Yes
AD Leo	10.01.2005	53381.51272	816	732	162.6476	2.1548E+32	0.519	Fast	Yes
AD Leo	10.01.2005	53381.52939	288	216	31.5306	4.1772E+31	0.399	Slow	Yes
AD Leo	10.02.2005	53412.49006	1308	1008	491.1727	6.5071E+32	0.863	Slow	No
AD Leo	10.02.2005	53412.52798	4164	3756	3907.4481	5.1767E+33	1.589	Fast	No
AD Leo	10.02.2005	53412.57464	1224	948	845.6318	1.1203E+33	0.792	Slow	No
AD Leo	10.02.2005	53412.58687	1764	1656	3250.0883	4.3058E+33	2.387	Fast	Yes
AD Leo	11.02.2005	53413.46661	36	24	3.0958	4.1014E+30	0.200	Slow	No
AD Leo	11.02.2005	53413.55759	384	360	60.6736	8.0382E+31	0.363	Fast	Yes
AD Leo	12.03.2005	53442.39045	720	680	78.4325	1.0391E+32	0.437	Fast	Yes
AD Leo	14.03.2005	53444.39896	432	300	30.9237	4.0968E+31	0.183	Slow	No
AD Leo	14.03.2005	53444.41327	1140	1068	152.7276	2.0234E+32	0.183	Fast	Yes
AD Leo	14.03.2005	53444.42674	588	504	54.4535	7.2141E+31	0.155	Fast	Yes
AD Leo	14.03.2005	53444.47105	1488	1428	320.0239	4.2397E+32	1.168	Fast	Yes
AD Leo	14.03.2005	53444.50563	1428	1356	206.4180	2.7347E+32	0.542	Fast	Yes
AD Leo	14.03.2005	53444.53049	240	72	25.4230	3.3681E+31	0.258	Slow	No
AD Leo	14.03.2005	53444.54716	576	324	47.9470	6.3521E+31	0.247	Slow	No
AD Leo	14.03.2005	53444.55396	84	60	10.7092	1.4188E+31	0.222	Slow	Yes
AD Leo	14.03.2005	53444.55507	84	48	5.5891	7.4046E+30	0.243	Slow	Yes
AD Leo	14.03.2005	53444.55632	132	72	8.6651	1.1480E+31	0.192	Slow	No
AD Leo	14.03.2005	53444.55966	24	12	8.6411	1.1448E+31	0.213	Slow	No
AD Leo	14.03.2005	53444.56021	60	36	8.6411	1.1448E+31	0.192	Slow	Yes

Table 3—Continued

Stars	Observation Date	HJD For Maximum of Flare	Total Time (sec)	Decay Time (sec)	Equivalent Duration (sec)	Flare Energy (ergs)	Amplitude In U Band (mag)	Flare Type	Used In Analyses
AD Leo	14.03.2005	53444.56313	204	156	24.4442	3.2384E+31	0.207	Slow	Yes
AD Leo	16.03.2005	53446.29405	2472	1812	293.6454	3.8903E+32	0.221	Slow	No
AD Leo	16.03.2005	53446.34849	48	24	2.4130	3.1968E+30	0.206	Slow	Yes
AD Leo	16.03.2005	53446.36113	456	408	33.4773	4.4351E+31	0.195	Fast	Yes
AD Leo	16.03.2005	53446.39558	2244	1032	198.8119	2.6339E+32	0.193	Slow	No
AD Leo	16.03.2005	53446.41252	852	756	229.8011	3.0444E+32	0.553	Fast	No
AD Leo	16.03.2005	53446.42960	756	588	99.5567	1.3189E+32	0.359	Slow	No
AD Leo	16.03.2005	53446.44822	132	72	12.2456	1.6223E+31	0.433	Slow	No
AD Leo	09.04.2005	53470.31255	1176	924	677.9344	8.9814E+32	1.259	Fast	No
AD Leo	09.04.2005	53470.33269	2088	1272	194.9401	2.5826E+32	0.121	Slow	No
AD Leo	09.04.2005	53470.36824	936	804	226.5383	3.0012E+32	0.938	Fast	No
AD Leo	10.04.2005	53471.30082	240	192	19.8520	2.6300E+31	0.234	Fast	Yes
AD Leo	10.04.2005	53471.37263	756	588	124.5185	1.6496E+32	0.512	Slow	No
AD Leo	02.05.2005	53493.31977	1284	1212	203.5007	2.6960E+32	0.699	Fast	Yes
AD Leo	09.05.2005	53500.32863	264	156	26.3800	3.4949E+31	0.207	Slow	No
AD Leo	09.05.2005	53500.35640	852	300	47.2850	6.2644E+31	0.164	Slow	No
AD Leo	08.01.2006	53744.50793	105	45	11.0325	1.4616E+31	0.236	Slow	No
AD Leo	08.01.2006	53744.55585	45	15	3.4390	4.5561E+30	0.162	Slow	No
AD Leo	08.01.2006	53744.64085	525	255	44.7429	5.9276E+31	0.192	Slow	No
AD Leo	27.01.2006	53763.62811	168	120	14.9402	1.9793E+31	0.171	Slow	Yes
AD Leo	27.01.2006	53763.63019	96	36	8.4330	1.1172E+31	0.162	Slow	No
AD Leo	02.02.2006	53769.52752	420	396	62.8336	8.3243E+31	0.510	Fast	Yes
AD Leo	02.02.2006	53769.55037	108	60	5.6936	7.5429E+30	0.140	Slow	Yes

Table 3—Continued

Stars	Observation Date	HJD For Maximum of Flare	Total Time (sec)	Decay Time (sec)	Equivalent Duration (sec)	Flare Energy (ergs)	Amplitude In U Band (mag)	Flare Type	Used In Analyses
AD Leo	02.02.2006	53769.61581	156	132	13.7184	1.8174E+31	0.138	Fast	Yes
AD Leo	02.02.2006	53769.61803	48	24	2.0431	2.7067E+30	0.161	Slow	Yes
AD Leo	02.02.2006	53769.62335	72	48	5.7062	7.5597E+30	0.160	Slow	Yes
AD Leo	02.02.2006	53769.62765	96	48	8.1257	1.0765E+31	0.173	Slow	Yes
AD Leo	04.02.2006	53771.46932	1188	1032	798.6289	1.0580E+33	1.210	Fast	No
AD Leo	04.02.2006	53771.49590	168	84	17.2768	2.2889E+31	0.321	Slow	Yes
AD Leo	04.02.2006	53771.50608	60	24	3.8830	5.1443E+30	0.183	Slow	Yes
AD Leo	21.02.2006	53788.42126	210	198	56.5353	7.4899E+31	0.618	Fast	No
AD Leo	21.02.2006	53788.42609	12	6	1.8036	2.3895E+30	0.261	Slow	No
AD Leo	21.02.2006	53788.49379	58	44	22.5737	2.9906E+31	0.741	Slow	Yes
AD Leo	21.02.2006	53788.49497	148	114	37.6468	4.9875E+31	0.522	Slow	Yes
AD Leo	21.02.2006	53788.49696	10	6	1.4739	1.9527E+30	0.290	Slow	No
AD Leo	21.02.2006	53788.53818	374	358	186.3927	2.4694E+32	1.365	Fast	Yes
AD Leo	21.02.2006	53788.54429	8	2	1.2042	1.5953E+30	0.267	Slow	No
AD Leo	21.02.2006	53788.54697	16	2	2.2268	2.9501E+30	0.337	Slow	Yes
AD Leo	17.03.2006	53812.29784	585	555	52.7018	6.9820E+31	0.294	Fast	Yes
AD Leo	17.03.2006	53812.32348	165	135	21.5461	2.8545E+31	0.230	Fast	Yes
AD Leo	25.03.2006	53820.28531	12	4	3.0728	4.0708E+30	0.363	Slow	No
AD Leo	25.03.2006	53820.31442	138	62	23.2816	3.0844E+31	0.358	Slow	No
AD Leo	25.03.2006	53820.31970	14	2	2.8248	3.7423E+30	0.582	Slow	No
AD Leo	25.03.2006	53820.40075	256	222	237.9794	3.1528E+32	1.171	Fast	Yes
AD Leo	25.03.2006	53820.40674	90	48	16.2012	2.1464E+31	0.435	Slow	No
AD Leo	25.03.2006	53820.40742	20	10	3.1643	4.1921E+30	0.542	Slow	No

Table 3—Continued

Stars	Observation Date	HJD For Maximum of Flare	Total Time (sec)	Decay Time (sec)	Equivalent Duration (sec)	Flare Energy (ergs)	Amplitude In U Band (mag)	Flare Type	Used In Analyses
AD Leo	25.03.2006	53820.41637	8	4	1.7195	2.2781E+30	0.448	Slow	No
AD Leo	25.03.2006	53820.41686	12	6	3.0961	4.1018E+30	0.477	Slow	No
AD Leo	01.04.2006	53827.27390	10	6	1.3930	1.8455E+30	0.464	Slow	No
AD Leo	01.04.2006	53827.29613	12	4	2.5516	3.3804E+30	0.493	Slow	No
AD Leo	01.04.2006	53827.30144	10	4	2.0011	2.6511E+30	0.461	Slow	No
AD Leo	01.04.2006	53827.31787	10	2	1.8332	2.4286E+30	0.519	Slow	No
AD Leo	01.04.2006	53827.37920	12	4	1.8984	2.5150E+30	0.427	Slow	No
AD Leo	05.04.2006	53831.39559	66	46	34.7529	4.6041E+31	0.713	Slow	Yes
AD Leo	05.04.2006	53831.39755	10	6	1.6828	2.2295E+30	0.446	Slow	No
AD Leo	05.04.2006	53831.40919	76	62	38.0426	5.0400E+31	0.917	Fast	No
AD Leo	05.04.2006	53831.41058	16	8	2.7778	3.6801E+30	0.459	Slow	No
AD Leo	05.04.2006	53831.41937	14	2	3.3755	4.4720E+30	0.536	Slow	No
AD Leo	01.12.2006	54071.52083	78	39	8.0437	1.0657E+31	0.190	Slow	No
AD Leo	01.12.2006	54071.52940	156	104	12.6557	1.6767E+31	0.229	Slow	No
AD Leo	01.12.2006	54071.53617	208	130	23.8354	3.1578E+31	0.208	Slow	No
AD Leo	01.12.2006	54071.54174	52	26	7.0330	9.3175E+30	0.186	Slow	No
AD Leo	01.12.2006	54071.59966	2258	1699	214.7257	2.8447E+32	0.201	Slow	No
AD Leo	15.12.2006	54085.63076	20	10	1.8367	2.4332E+30	0.139	Slow	No
AD Leo	15.12.2006	54085.63572	30	20	1.9239	2.5488E+30	0.160	Slow	No
AD Leo	23.12.2006	54093.60562	30	10	3.2230	4.2698E+30	0.151	Slow	Yes
AD Leo	23.12.2006	54093.60712	40	10	3.5123	4.6532E+30	0.170	Slow	No
AD Leo	23.12.2006	54093.61349	150	110	10.6923	1.4165E+31	0.195	Slow	No
AD Leo	23.12.2006	54093.63486	650	400	58.1135	7.6990E+31	0.186	Slow	No

Table 3—Continued

Stars	Observation Date	HJD For Maximum of Flare	Total Time (sec)	Decay Time (sec)	Equivalent Duration (sec)	Flare Energy (ergs)	Amplitude In U Band (mag)	Flare Type	Used In Analyses
AD Leo	23.12.2006	54093.64631	1676	1086	154.4312	2.0459E+32	0.200	Slow	No
AD Leo	21.01.2007	54122.47422	16	8	1.8714	2.4792E+30	0.229	Slow	No
AD Leo	21.01.2007	54122.48635	32	16	3.9082	5.1777E+30	0.274	Slow	Yes
AD Leo	21.01.2007	54122.48876	16	8	2.0181	2.6736E+30	0.302	Slow	No
AD Leo	21.01.2007	54122.49107	32	16	3.6712	4.8637E+30	0.213	Slow	Yes
AD Leo	21.01.2007	54122.49573	24	16	2.9831	3.9520E+30	0.248	Slow	No
AD Leo	21.01.2007	54122.49693	16	8	1.7260	2.2867E+30	0.264	Slow	No
AD Leo	21.01.2007	54122.49712	16	8	1.7700	2.3449E+30	0.215	Slow	No
AD Leo	21.01.2007	54122.49860	16	8	1.8787	2.4889E+30	0.214	Slow	No
AD Leo	21.01.2007	54122.50156	24	8	2.9281	3.8792E+30	0.259	Slow	Yes
AD Leo	21.01.2007	54122.50582	40	24	3.0146	3.9938E+30	0.211	Slow	Yes
AD Leo	21.01.2007	54122.50628	24	16	2.5792	3.4170E+30	0.217	Slow	No
AD Leo	21.01.2007	54122.53384	24	16	3.2242	4.2715E+30	0.279	Slow	No
AD Leo	08.03.2007	54168.46811	60	30	8.2226	1.0893E+31	0.285	Slow	No
AD Leo	08.03.2007	54168.46933	90	60	14.2830	1.8922E+31	0.232	Slow	No
AD Leo	16.03.2007	54176.37237	84	60	4.8172	6.3820E+30	0.157	Slow	Yes
AD Leo	16.03.2007	54176.39611	192	120	18.7237	2.4805E+31	0.230	Slow	Yes
AD Leo	16.03.2007	54176.45538	372	156	28.2549	3.7433E+31	0.168	Slow	No
EQ Peg	18.08.2004	53236.43176	290	250	44.8869	4.4695E+31	0.369	Fast	Yes
EQ Peg	18.08.2004	53236.48106	210	170	28.0058	2.7886E+31	0.226	Fast	Yes
EQ Peg	18.08.2004	53236.50398	220	180	42.7730	4.2590E+31	0.473	Fast	Yes
EQ Peg	18.08.2004	53236.56139	1060	900	475.3540	4.7332E+32	1.279	Fast	No
EQ Peg	19.08.2004	53237.38157	490	440	111.7547	1.1128E+32	0.503	Fast	Yes

Table 3—Continued

Stars	Observation Date	HJD For Maximum of Flare	Total Time (sec)	Decay Time (sec)	Equivalent Duration (sec)	Flare Energy (ergs)	Amplitude In U Band (mag)	Flare Type	Used In Analyses
EQ Peg	19.08.2004	53237.40715	150	110	34.0142	3.3869E+31	0.258	Slow	No
EQ Peg	19.08.2004	53237.41224	340	310	68.8944	6.8600E+31	0.464	Fast	Yes
EQ Peg	19.08.2004	53237.48805	230	200	51.5306	5.1310E+31	0.734	Fast	Yes
EQ Peg	19.08.2004	53237.49291	180	160	30.2882	3.0159E+31	0.343	Fast	Yes
EQ Peg	19.08.2004	53237.53122	610	380	106.3842	1.0593E+32	0.384	Slow	No
EQ Peg	22.08.2004	53240.36934	570	510	260.2007	2.5909E+32	0.880	Fast	Yes
EQ Peg	08.09.2004	53257.36799	340	270	56.6293	5.6387E+31	0.427	Fast	No
EQ Peg	08.09.2004	53257.38223	430	140	72.0350	7.1727E+31	0.478	Slow	No
EQ Peg	08.09.2004	53257.45202	1170	1040	199.7623	1.9891E+32	0.406	Fast	No
EQ Peg	08.09.2004	53257.51000	120	110	14.8846	1.4821E+31	0.413	Fast	No
EQ Peg	08.09.2004	53257.51845	60	30	11.3037	1.1255E+31	0.520	Slow	No
EQ Peg	09.09.2004	53258.36870	1840	1390	409.0358	4.0729E+32	0.478	Slow	No
EQ Peg	09.09.2004	53258.40585	190	160	32.7460	3.2606E+31	0.574	Fast	Yes
EQ Peg	09.09.2004	53258.51094	200	160	33.3344	3.3192E+31	0.484	Fast	Yes
EQ Peg	12.09.2004	53261.39106	260	220	53.8905	5.3660E+31	0.583	Fast	Yes
EQ Peg	12.09.2004	53261.46988	130	110	19.8878	1.9803E+31	0.332	Fast	Yes
EQ Peg	12.09.2004	53261.55529	150	130	35.5859	3.5434E+31	0.756	Fast	Yes
EQ Peg	12.09.2004	53261.59048	210	180	74.9052	7.4585E+31	0.785	Fast	Yes
EQ Peg	14.09.2004	53263.34072	1070	950	110.0519	1.0958E+32	0.251	Fast	No
EQ Peg	14.09.2004	53263.36815	190	130	41.2794	4.1103E+31	0.532	Slow	No
EQ Peg	14.09.2004	53263.37752	1620	1430	625.0200	6.2235E+32	0.868	Fast	No
EQ Peg	14.09.2004	53263.40484	200	150	44.5257	4.4335E+31	0.414	Slow	Yes
EQ Peg	14.09.2004	53263.47371	50	40	4.9279	4.9068E+30	0.289	Fast	No

Table 3—Continued

Stars	Observation Date	HJD For Maximum of Flare	Total Time (sec)	Decay Time (sec)	Equivalent Duration (sec)	Flare Energy (ergs)	Amplitude In U Band (mag)	Flare Type	Used In Analyses
EQ Peg	14.09.2004	53263.48725	70	50	9.4754	9.4349E+30	0.357	Slow	Yes
EQ Peg	14.09.2004	53263.54697	60	30	5.7809	5.7562E+30	0.361	Slow	No
EQ Peg	15.09.2004	53264.46607	330	120	44.1024	4.3914E+31	0.246	Slow	No
EQ Peg	15.09.2004	53264.50102	580	550	294.4328	2.9317E+32	2.109	Fast	Yes
EQ Peg	15.09.2004	53264.51167	510	490	166.0980	1.6539E+32	1.420	Fast	Yes
EQ Peg	15.09.2004	53264.56595	1440	390	186.4048	1.8561E+32	0.284	Slow	No
EQ Peg	16.09.2004	53265.39674	910	860	180.5574	1.7979E+32	0.622	Fast	Yes
EQ Peg	16.09.2004	53265.51329	420	250	84.7088	8.4347E+31	0.203	Slow	No
EQ Peg	16.09.2004	53265.54003	410	350	99.4698	9.9044E+31	0.463	Fast	Yes
EQ Peg	16.09.2004	53265.56942	3180	1950	746.5378	7.4335E+32	0.345	Slow	No
EQ Peg	07.09.2005	53621.52654	150	105	34.9365	3.4787E+31	0.630	Slow	No
EQ Peg	08.09.2005	53622.45159	285	270	78.7774	7.8441E+31	0.711	Fast	Yes
EQ Peg	08.09.2005	53622.46218	540	510	156.1378	1.5547E+32	0.968	Fast	Yes
EQ Peg	08.09.2005	53622.48087	120	105	39.0373	3.8870E+31	0.640	Fast	Yes
EQ Peg	08.09.2005	53622.49927	75	60	12.4900	1.2437E+31	0.362	Fast	Yes
EQ Peg	12.09.2005	53626.33898	1839	1764	6818.1425	6.7890E+33	4.006	Fast	Yes
EQ Peg	12.09.2005	53626.36062	75	45	8.6906	8.6534E+30	0.261	Slow	No
EQ Peg	12.09.2005	53626.36218	105	75	15.2364	1.5171E+31	0.338	Slow	No
EQ Peg	12.09.2005	53626.36409	105	30	10.5085	1.0464E+31	0.337	Slow	Yes
EQ Peg	12.09.2005	53626.36582	922	802	352.2107	3.5070E+32	1.285	Fast	No
EQ Peg	12.09.2005	53626.40874	225	180	51.6810	5.1460E+31	0.567	Fast	Yes
EQ Peg	12.09.2005	53626.50223	90	45	24.2883	2.4184E+31	0.440	Slow	No
EQ Peg	12.09.2005	53626.56201	315	225	181.2224	1.8045E+32	0.964	Slow	Yes

Table 3—Continued

Stars	Observation Date	HJD For Maximum of Flare	Total Time (sec)	Decay Time (sec)	Equivalent Duration (sec)	Flare Energy (ergs)	Amplitude In U Band (mag)	Flare Type	Used In Analyses
EQ Peg	12.09.2005	53626.56478	135	120	83.2370	8.2881E+31	1.090	Fast	Yes
EQ Peg	12.09.2005	53626.56635	60	45	12.1029	1.2051E+31	0.252	Slow	Yes
EQ Peg	12.09.2005	53626.56721	180	150	27.7286	2.7610E+31	0.282	Fast	Yes
EQ Peg	12.09.2005	53626.56930	60	30	5.7164	5.6920E+30	0.248	Slow	No
EQ Peg	12.09.2005	53626.58426	165	135	31.8950	3.1759E+31	0.676	Fast	Yes
EQ Peg	27.09.2005	53641.34719	135	120	76.2420	7.5916E+31	1.111	Fast	Yes
EQ Peg	27.09.2005	53641.36437	90	45	9.0307	8.9921E+30	0.225	Slow	No
EQ Peg	27.09.2005	53641.39000	150	120	35.8623	3.5709E+31	0.535	Fast	Yes
EQ Peg	27.09.2005	53641.42135	1545	1410	1739.8937	1.7325E+33	1.762	Fast	No
EQ Peg	27.09.2005	53641.47316	977	585	203.3585	2.0249E+32	0.387	Slow	No
EQ Peg	03.10.2005	53647.34180	180	135	66.8966	6.6611E+31	1.076	Slow	No
EQ Peg	03.10.2005	53647.40952	255	180	56.0173	5.5778E+31	0.536	Slow	Yes
EQ Peg	28.10.2005	53672.27432	2764	1594	378.4947	3.7688E+32	0.333	Slow	No
EQ Peg	28.10.2005	53672.30797	210	120	52.2187	5.1995E+31	0.402	Slow	Yes
EQ Peg	28.10.2005	53672.33656	225	195	55.9946	5.5755E+31	0.682	Fast	Yes
EQ Peg	29.10.2005	53673.34763	570	450	155.7805	1.5511E+32	0.740	Fast	No
EQ Peg	29.10.2005	53673.35896	240	135	72.3348	7.2025E+31	0.716	Slow	No
EQ Peg	29.10.2005	53673.36104	330	285	96.7821	9.6368E+31	0.461	Fast	Yes
EQ Peg	29.10.2005	53673.38314	738	468	171.8148	1.7108E+32	0.428	Slow	No
EQ Peg	11.11.2005	53686.29364	540	405	151.3632	1.5072E+32	0.511	Slow	No
EQ Peg	11.11.2005	53686.30076	795	645	566.9639	5.6454E+32	1.019	Fast	No
EQ Peg	11.11.2005	53686.37753	270	120	97.9336	9.7515E+31	0.562	Slow	No
EV Lac	11.07.2004	53198.47514	260	190	50.2493	2.3691E+31	0.431	Slow	No

Table 3—Continued

Stars	Observation Date	HJD For Maximum of Flare	Total Time (sec)	Decay Time (sec)	Equivalent Duration (sec)	Flare Energy (ergs)	Amplitude In U Band (mag)	Flare Type	Used In Analyses
EV Lac	11.07.2004	53198.48325	590	560	128.0124	6.0353E+31	0.735	Fast	Yes
EV Lac	17.07.2004	53204.51795	1230	820	174.6712	8.2351E+31	0.313	Slow	No
EV Lac	20.07.2004	53207.49970	310	230	64.2165	3.0276E+31	0.588	Slow	No
EV Lac	20.07.2004	53207.51648	510	490	114.2672	5.3873E+31	0.905	Fast	Yes
EV Lac	24.07.2004	53211.54445	830	590	165.8313	7.8184E+31	0.397	Slow	No
EV Lac	25.07.2004	53212.48570	490	340	141.3253	6.6630E+31	0.602	Slow	No
EV Lac	25.07.2004	53212.51625	380	160	167.7941	7.9109E+31	0.514	Slow	No
EV Lac	26.07.2004	53213.48702	80	50	17.0178	8.0233E+30	0.676	Slow	No
EV Lac	26.07.2004	53213.52209	100	90	15.4025	7.2618E+30	0.385	Fast	No
EV Lac	28.07.2004	53215.48409	180	160	97.5996	4.6015E+31	1.315	Fast	Yes
EV Lac	28.07.2004	53215.49960	170	120	36.9820	1.7436E+31	0.436	Slow	Yes
EV Lac	07.08.2004	53225.42498	350	190	94.2676	4.4444E+31	0.611	Slow	No
EV Lac	07.08.2004	53225.46827	140	80	47.9538	2.2609E+31	0.643	Slow	No
EV Lac	08.08.2004	53226.46483	140	120	26.0719	1.2292E+31	0.377	Fast	Yes
EV Lac	08.08.2004	53226.50951	2482	2422	594.8340	2.8044E+32	0.816	Fast	Yes
EV Lac	09.08.2004	53227.38165	940	910	728.2759	3.4336E+32	1.763	Fast	Yes
EV Lac	09.08.2004	53227.42505	630	490	188.9706	8.9093E+31	0.555	Slow	No
EV Lac	09.08.2004	53227.43928	360	260	83.5153	3.9375E+31	0.591	Slow	No
EV Lac	09.08.2004	53227.44287	240	220	44.6927	2.1071E+31	0.427	Fast	Yes
EV Lac	09.08.2004	53227.46382	1000	960	1134.4908	5.3487E+32	1.987	Fast	Yes
EV Lac	09.08.2004	53227.49692	90	70	18.4229	8.6857E+30	0.511	Slow	Yes
EV Lac	09.08.2004	53227.55063	70	60	9.8307	4.6348E+30	0.353	Fast	No
EV Lac	10.08.2004	53228.37056	90	50	19.5669	9.2251E+30	0.404	Slow	No

Table 3—Continued

Stars	Observation Date	HJD For Maximum of Flare	Total Time (sec)	Decay Time (sec)	Equivalent Duration (sec)	Flare Energy (ergs)	Amplitude In U Band (mag)	Flare Type	Used In Analyses
EV Lac	10.08.2004	53228.38654	2200	2090	4235.9481	1.9971E+33	2.718	Fast	No
EV Lac	10.08.2004	53228.41698	1800	1730	849.9341	4.0071E+32	2.020	Fast	No
EV Lac	10.08.2004	53228.52716	50	20	8.2244	3.8775E+30	0.437	Slow	No
EV Lac	10.08.2004	53228.55471	120	100	24.2433	1.1430E+31	0.520	Fast	Yes
EV Lac	14.08.2004	53232.36686	80	50	14.2793	6.7322E+30	0.433	Slow	No
EV Lac	14.08.2004	53232.37785	290	220	72.8660	3.4354E+31	0.660	Slow	No
EV Lac	14.08.2004	53232.39359	520	470	147.9080	6.9733E+31	0.751	Fast	Yes
EV Lac	05.07.2005	53557.52397	168	116	71.6667	3.3788E+31	0.693	Slow	No
EV Lac	05.07.2005	53557.53656	4	2	1.4724	6.9419E+29	0.516	Slow	No
EV Lac	26.07.2005	53578.50257	600	585	184.9502	8.7197E+31	0.727	Fast	Yes
EV Lac	26.07.2005	53578.51594	105	75	15.5441	7.3285E+30	0.381	Slow	No
EV Lac	26.07.2005	53578.52566	30	15	3.7640	1.7746E+30	0.381	Slow	Yes
EV Lac	26.07.2005	53578.54667	120	45	12.2619	5.7810E+30	0.372	Slow	Yes
EV Lac	03.08.2005	53586.50688	570	480	384.1497	1.8111E+32	1.565	Fast	Yes
EV Lac	03.08.2005	53586.51348	30	15	2.6479	1.2484E+30	0.313	Slow	Yes
EV Lac	03.08.2005	53586.52856	180	135	38.1427	1.7983E+31	0.437	Slow	No
EV Lac	14.08.2005	53597.33755	135	105	41.7339	1.9676E+31	0.361	Slow	No
EV Lac	14.08.2005	53597.40029	60	45	8.4471	3.9825E+30	0.297	Slow	Yes
EV Lac	14.08.2005	53597.47199	105	60	23.3056	1.0988E+31	0.359	Slow	No
EV Lac	14.08.2005	53597.47615	210	120	44.6968	2.1073E+31	0.401	Slow	Yes
EV Lac	14.08.2005	53597.47806	105	75	35.7253	1.6843E+31	0.304	Slow	No
EV Lac	14.08.2005	53597.48015	105	75	30.4018	1.4333E+31	0.493	Slow	No
EV Lac	14.08.2005	53597.48223	165	60	32.2519	1.5206E+31	0.228	Slow	No

Table 3—Continued

Stars	Observation Date	HJD For Maximum of Flare	Total Time (sec)	Decay Time (sec)	Equivalent Duration (sec)	Flare Energy (ergs)	Amplitude In U Band (mag)	Flare Type	Used In Analyses
EV Lac	15.08.2005	53598.42911	390	345	92.8472	4.3774E+31	0.435	Fast	Yes
EV Lac	15.08.2005	53598.43380	195	165	33.7418	1.5908E+31	0.219	Fast	Yes
EV Lac	15.08.2005	53598.46517	1005	870	398.3372	1.8780E+32	0.949	Fast	No
EV Lac	15.08.2005	53598.50008	810	720	282.6663	1.3327E+32	1.023	Fast	Yes
EV Lac	15.08.2005	53598.52741	1365	1245	1264.2000	5.9603E+32	2.096	Fast	No
EV Lac	22.08.2005	53605.39060	60	45	21.6520	1.0208E+31	0.467	Slow	Yes
EV Lac	22.08.2005	53605.39147	330	300	187.0415	8.8183E+31	1.058	Fast	Yes
EV Lac	22.08.2005	53605.41460	195	105	61.7938	2.9134E+31	0.469	Slow	Yes
EV Lac	22.08.2005	53605.42241	75	30	20.6508	9.7361E+30	0.496	Slow	No
EV Lac	22.08.2005	53605.42467	105	75	24.9765	1.1776E+31	0.439	Slow	No
EV Lac	22.08.2005	53605.43113	45	30	12.4383	5.8642E+30	0.434	Slow	Yes
EV Lac	23.08.2005	53606.38421	45	30	8.4172	3.9684E+30	0.312	Slow	Yes
EV Lac	23.08.2005	53606.42194	45	30	7.6081	3.5869E+30	0.460	Slow	Yes
EV Lac	23.08.2005	53606.45404	120	105	30.8166	1.4529E+31	0.333	Fast	Yes
EV Lac	23.08.2005	53606.49226	45	15	1.6396	7.7299E+29	0.476	Slow	No
EV Lac	23.08.2005	53606.49851	30	15	3.5732	1.6846E+30	0.412	Slow	Yes
EV Lac	23.07.2006	53940.47444	60	30	7.9475	3.7470E+30	0.422	Slow	No
EV Lac	23.07.2006	53940.52270	480	375	113.1827	5.3362E+31	0.572	Fast	No
EV Lac	23.07.2006	53940.53726	120	90	16.4250	7.7438E+30	0.334	Slow	No
EV Lac	29.07.2006	53946.47599	1054	989	570.7213	2.6907E+32	1.991	Fast	No
EV Lac	29.07.2006	53946.54317	75	45	12.2062	5.7548E+30	0.394	Slow	No
EV Lac	03.08.2006	53951.53751	2898	2688	2579.1462	1.2160E+33	1.574	Fast	No
EV Lac	04.08.2006	53952.37326	125	65	12.4023	5.8473E+30	0.238	Slow	No

Table 3—Continued

Stars	Observation Date	HJD For Maximum of Flare	Total Time (sec)	Decay Time (sec)	Equivalent Duration (sec)	Flare Energy (ergs)	Amplitude In U Band (mag)	Flare Type	Used In Analyses
EV Lac	04.08.2006	53952.37413	30	20	3.6127	1.7033E+30	0.334	Slow	No
EV Lac	04.08.2006	53952.37911	20	10	2.8304	1.3344E+30	0.314	Slow	No
EV Lac	04.08.2006	53952.37934	20	10	2.6850	1.2659E+30	0.345	Slow	No
EV Lac	04.08.2006	53952.37980	30	20	4.2637	2.0102E+30	0.319	Slow	No
EV Lac	07.08.2006	53955.48708	90	60	14.2999	6.7419E+30	0.460	Slow	No
EV Lac	07.08.2006	53955.52215	90	60	7.6945	3.6277E+30	0.480	Slow	No
EV Lac	07.08.2006	53955.53095	120	90	28.0671	1.3233E+31	0.578	Slow	No
EV Lac	08.08.2006	53956.34904	60	45	13.7110	6.4642E+30	0.652	Slow	Yes
EV Lac	08.08.2006	53956.35441	120	105	38.2660	1.8041E+31	0.633	Fast	Yes
EV Lac	08.08.2006	53956.39257	1680	1590	900.4373	4.2452E+32	1.484	Fast	Yes
EV Lac	08.08.2006	53956.44677	30	15	8.5613	4.0364E+30	0.459	Slow	Yes
EV Lac	08.08.2006	53956.44989	150	75	23.5724	1.1114E+31	0.489	Slow	Yes
EV Lac	08.08.2006	53956.45215	75	45	16.8672	7.9523E+30	0.516	Slow	No
EV Lac	08.08.2006	53956.45285	30	15	8.7699	4.1347E+30	0.450	Slow	Yes
EV Lac	08.08.2006	53956.45510	195	105	56.8137	2.6786E+31	0.486	Slow	Yes
EV Lac	12.08.2006	53960.53098	2940	2100	1453.1544	6.8511E+32	0.821	Slow	No
EV Lac	15.08.2006	53963.40072	30	10	7.4410	3.5082E+30	0.578	Slow	Yes
EV Lac	25.08.2006	53973.49652	2330	2260	2288.6227	1.0790E+33	2.164	Fast	No
EV Lac	05.09.2006	53984.44139	360	315	253.0786	1.1932E+32	1.215	Fast	Yes
EV Lac	05.09.2006	53984.47750	135	60	35.1887	1.6590E+31	0.563	Slow	Yes
EV Lac	05.09.2006	53984.50997	30	15	8.6037	4.0564E+30	0.475	Slow	Yes
EV Lac	08.09.2006	53987.47180	90	45	19.8145	9.3418E+30	0.537	Slow	No
EV Lac	08.09.2006	53987.47389	75	30	24.6323	1.1613E+31	0.845	Slow	No

Table 3—Continued

Stars	Observation Date	HJD For Maximum of Flare	Total Time (sec)	Decay Time (sec)	Equivalent Duration (sec)	Flare Energy (ergs)	Amplitude In U Band (mag)	Flare Type	Used In Analyses
EV Lac	15.09.2006	53994.36716	70	20	19.0094	8.9622E+30	0.441	Slow	Yes
EV Lac	15.09.2006	53994.37087	320	150	59.0751	2.7852E+31	0.359	Slow	No
EV Lac	17.09.2006	53996.33563	435	390	82.6432	3.8963E+31	0.395	Fast	Yes
EV Lac	17.09.2006	53996.34258	360	150	65.9648	3.1100E+31	0.253	Slow	No
EV Lac	17.09.2006	53996.35595	195	135	35.6703	1.6817E+31	0.317	Slow	No
V1054 Oph	11.06.2004	53168.42510	140	80	13.5776	3.9221E+31	0.274	Slow	No
V1054 Oph	14.06.2004	53171.39288	650	600	440.3598	1.2720E+33	1.829	Fast	Yes
V1054 Oph	14.06.2004	53171.42147	180	130	10.7079	3.0931E+31	0.226	Slow	Yes
V1054 Oph	14.06.2004	53171.42563	3270	3190	1441.3629	4.1636E+33	1.239	Fast	Yes
V1054 Oph	14.06.2004	53171.46290	1780	1750	281.5233	8.1322E+32	0.359	Fast	Yes
V1054 Oph	20.06.2004	53177.40665	300	270	32.6526	9.4321E+31	0.355	Fast	Yes
V1054 Oph	20.06.2004	53177.42459	120	60	13.8765	4.0084E+31	0.230	Slow	No
V1054 Oph	20.06.2004	53177.43755	1000	520	59.0017	1.7043E+32	0.181	Slow	No
V1054 Oph	04.07.2004	53191.36033	340	90	43.0587	1.2438E+32	0.305	Slow	No
V1054 Oph	04.07.2004	53191.36334	1340	1280	329.4567	9.5168E+32	0.957	Fast	Yes
V1054 Oph	04.07.2004	53191.39505	1800	340	247.6494	7.1537E+32	0.194	Slow	No
V1054 Oph	06.07.2004	53193.34635	1070	960	429.5400	1.2408E+33	1.568	Fast	No
V1054 Oph	06.07.2004	53193.36302	1460	980	192.7904	5.5690E+32	0.180	Slow	No
V1054 Oph	10.07.2004	53197.39384	670	530	88.1072	2.5451E+32	0.502	Fast	No
V1054 Oph	11.05.2005	53502.41191	48	36	2.3143	6.6853E+30	0.151	Slow	No
V1054 Oph	11.05.2005	53502.41469	216	180	24.2944	7.0178E+31	0.365	Fast	Yes
V1054 Oph	04.06.2005	53526.42970	36	24	2.2093	6.3819E+30	0.122	Slow	No
V1054 Oph	04.06.2005	53526.44400	60	36	6.1596	1.7793E+31	0.175	Slow	Yes

Table 3—Continued

Stars	Observation Date	HJD For Maximum of Flare	Total Time (sec)	Decay Time (sec)	Equivalent Duration (sec)	Flare Energy (ergs)	Amplitude In U Band (mag)	Flare Type	Used In Analyses
V1054 Oph	05.06.2005	53527.40053	468	396	51.9875	1.5017E+32	0.271	Fast	Yes
V1054 Oph	05.06.2005	53527.40636	48	36	3.9567	1.1429E+31	0.192	Slow	No
V1054 Oph	05.06.2005	53527.45803	108	84	5.6538	1.6332E+31	0.152	Slow	Yes
V1054 Oph	06.06.2005	53528.38594	1596	708	160.4190	4.6339E+32	0.250	Slow	No
V1054 Oph	06.06.2005	53528.39942	1368	1008	168.3597	4.8633E+32	0.207	Slow	No
V1054 Oph	06.06.2005	53528.41775	288	276	31.2711	9.0331E+31	0.227	Fast	No
V1054 Oph	13.06.2005	53535.44630	2304	1260	280.2476	8.0953E+32	0.258	Slow	No
V1054 Oph	13.06.2005	53535.47463	1416	768	102.0127	2.9468E+32	0.182	Slow	No
V1054 Oph	13.06.2005	53535.48505	192	156	12.1108	3.4984E+31	0.216	Fast	Yes
V1054 Oph	13.06.2005	53535.49797	144	96	11.4981	3.3214E+31	0.314	Slow	Yes
V1054 Oph	13.06.2005	53535.50880	132	120	18.6571	5.3893E+31	0.395	Fast	No
V1054 Oph	24.06.2005	53546.42356	72	48	8.2024	2.3694E+31	0.374	Slow	Yes
V1054 Oph	24.06.2005	53546.48564	48	36	2.6777	7.7348E+30	0.227	Slow	No
V1054 Oph	24.06.2005	53546.48717	48	36	5.5471	1.6023E+31	0.295	Slow	No
V1054 Oph	24.06.2005	53546.48912	36	24	3.2846	9.4881E+30	0.242	Slow	No
V1054 Oph	24.06.2005	53546.49148	36	24	3.4479	9.9596E+30	0.266	Slow	No
V1054 Oph	26.06.2005	53548.40336	48	36	1.5149	4.3761E+30	0.151	Slow	No
V1054 Oph	26.06.2005	53548.47419	36	12	4.9893	1.4412E+31	0.421	Slow	Yes
V1054 Oph	26.06.2005	53548.48919	300	276	10.4920	3.0308E+31	0.313	Fast	Yes
V1054 Oph	02.07.2005	53554.36689	108	60	5.4493	1.5741E+31	0.212	Slow	Yes
V1054 Oph	03.07.2005	53555.33448	564	36	33.8135	9.7675E+31	0.209	Slow	No
V1054 Oph	03.07.2005	53555.33921	960	624	103.7402	2.9967E+32	0.231	Slow	No

Table 4: For both fast and slow flares whose rise times are the same, the results obtained from both the regression calculations and the *t-Test* analyses performed to the mean averages of the equivalent durations ($\text{Log}(P_u)$) versus flare rise times ($\text{Log}(T_r)$) in logarithmic scale are listed.

Flare Groups :	Slow Flare	Fast Flare
<i>Data</i>		
Total Flare Number :	30	30
<i>Best Representation Values</i>		
Slope :	1.109 ± 0.127	1.227 ± 0.243
y intercept when $x = 0.0$:	-0.581 ± 0.226	0.122 ± 0.433
x intercept when $y = 0.0$:	0.524	-0.099
<i>Mean Average of All Y Values</i>		
Mean Average :	1.348	2.255
Mean Average Error :	0.092	0.126
<i>Goodness of Fit</i>		
r^2 :	0.732	0.476
<i>Is slope significantly non-zero?</i>		
p -value :	< 0.0001	< 0.0001
Deviation from zero? :	<i>Significant</i>	<i>Significant</i>